



# Department of Pesticide Regulation



Mary-Ann Warmerdam  
Director

Arnold Schwarzenegger  
Governor

March 28, 2007

ID #222113

Mr. Robert Prince  
President  
Microgen, Inc.  
33 Clinton Road, Suite 102  
Clinton Square Executive Center  
West Caldwell, NJ 07006

Dear Mr. Prince:

PUBLIC PLACES, EPA REG. NO. 61178-2-ML  
Label Amendments to Revise Precautionary and First Aid Statements  
and Revise Fungicidal and Deodorizer Directions

The above-amended label was registered effective March 28, 2007.

This registration action is based on compliance with provisions of California laws and regulations pertaining to pesticide registration. It is on file as the "latest" registered Master Label.

A stamped, accepted copy of the amended label is enclosed:

Sincerely,

Steve Rhodes  
Senior Pesticide Use Specialist  
Pesticide Registration Branch  
E-mail: [srhodes@cdpr.ca.gov](mailto:srhodes@cdpr.ca.gov)

Enclosure



**BOTTLE:FRONT PANEL**

**PUBLIC PLACES**

Virucidal\* Disinfectant spray, Cleans as it disinfects, Redefining Clean  
Ready to Use, Ready to Use Spray, Ready to Use Solution  
Contains Public Places Brand Hospital Disinfectant Formula Exclusively from Microgen, Inc

**ACTIVE INGREDIENTS**

Alkyl (C14 60%,C16 30%,C12 5%,C18 5%) dimethyl benzyl ammonium chloride....0.07%

Alkyl (C12 68%,C14 32%) dimethyl ethylbenzyl ammonium chloride.....0.07%

INERT INGREDIENTS.....99.86%

TOTAL.....100.00%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back panel for additional precautionary statements.

Net Contents: 1-32 ounces

**BOTTLE: BACK PANEL**

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

A one-step disinfectant spray that works in the presence of 5% organic load. For use on toilet seats, telephones, door knobs and similar hard nonporous inanimate environmental surfaces in homes, restaurants, hotels, motels, gas stations, health clubs and spas. Controls cross contamination from treated inanimate environmental surfaces.

**TO DISINFECT/ CLEAN/ DEODORIZE:**

Preclean all heavily soiled surfaces prior product application. To clean, deodorize, and disinfect hold dispenser 10 inches from the surface, press atomizer with quick short strokes, spraying evenly until surface is thoroughly wet. Wait 10 minutes, and wipe dry or air dry. For food contact surfaces such as tables and counters, rinse with potable water. Do not use on glasses, dishes or utensils

Refer to insert for complete organism contact times.

**PRECAUTIONARY STATEMENTS**

Hazard to Humans and Domestic Animals.

**Caution.** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or tobacco. Remove contaminated clothing and wash before reuse.

**FIRST AID**

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**Have the product container or label with you when calling a poison control center or doctor, or going for treatment.**

(For containers 1 gallon or less)

**STORAGE AND DISPOSAL**

Store in area inaccessible to children and person unfamiliar with product use. Do not reuse empty container. Discard in trash.


Microgen Inc.

33 Clinton Road Suite 102

West Caldwell, NJ 07006

EPA REG NO. 61178-2 EPA EST. NO.

CAL P/P (02/26/2007)



## **CARD- FRONT PANEL**

For pocket, purse, or institutional use

For travel or work

### **SPRAY ONTO:**

Toilet Seats

Telephones, Furniture

Door Knobs, Light Switches

Bathroom Fixtures, Sinks

Bathtubs, Shower Stalls

Stoves, Exterior Surfaces of Refrigerators

Interior Surfaces of Commercial Refrigerators\*\*

Counters, Tables

### **FOR USE IN:**

Public Restrooms

Airplanes

Hotels, Motels

Day Care Centers/Nursing Homes

Highway Service Areas/Gas Stations

Hospitals/Doctors-Dentist Offices & Labs/Clinics & Sickrooms

Veterinary Clinics

Health Clubs/Spas/Gyms

Restaurants/Bars/Fast Food Chains

Homes, Offices

Laundromats

Camp Sites, RV's

### **FOR USE BY:**

Vacationers

Business Travelers

Truckers

Doctors, Nurses

Paramedics

CPR Instructors

Firefighters, Police

\*\*Note to users in commercial settings: Interior surfaces of refrigerators are classified by the USEPA as a food contact surface and require the user to conduct a potable water rinse on the interior surfaces of the refrigerator after the interior surfaces are treated with this product.

**\*KILLS HIV-1 AND HBV ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV).

**SPECIAL INSTRUCTIONS FOR CLEANING & DECONTAMINATION AGAINST HIV-1 AND HBV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:**

**PERSONAL PROTECTION:** Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks, or eye coverings.

**CLEANING PROCEDURE:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the disinfectant.

**DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

**CONTACT TIME:** Leave surfaces wet for 10 minutes and 30 seconds for HBV and HIV-1, respectively. Use 10 minute contact time to mitigate other viruses, bacteria and fungi listed on the label and/or on the insert unless noted for specific organisms which require longer contact time.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

**SANITIZATION - NON-FOOD CONTACT SURFACES:** Remove all gross filth or heavy soil prior to applying sanitizing solution. To sanitize walls, floors and other hard, nonporous surfaces in such areas as schools, institutions, and industries, spray this solution so as to wet all surfaces thoroughly for 1 minute. Drain or air dry.

**MILDEWSTAT:** To control mold and mildew on pre-cleaned, hard, non-porous surfaces (such as floors, walls, table tops). Spray solution making sure to wet all surfaces completely. Let air dry. Repeat application at weekly intervals or when mildew growth reappears.

**DEODORIZER:** This product deodorizes garbage storage areas, empty garbage bins and cans, exterior surfaces of toilet bowls and any other odor-causing areas. Spray solution making sure to wet all surfaces completely. Allow to air dry.

**FUNGICIDAL:** Kills Trichophyton mentagrophytes, agent causes Athlete's Foot Fungus, on hard nonporous surfaces in areas such as locker rooms, exercise facilities and bath and shower areas. Spray solution making sure to wet all surfaces completely. Wait 10 minutes, then remove excess liquid or allow to air dry.

**FOR VETERINARY CLINICS DISINFECTANT USE**

1. Remove all animals and feed from premises, cars, boats, trucks and other equipment.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
6. Treat all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set or dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

**DISINFECTION OF FOOD PROCESSING AREAS:** Before use in food processing areas, all food products and packaging materials must be removed from the room or carefully protected. Spray solution making sure to wet all surfaces completely. Wait 10 minutes, then remove excess liquid or allow to air dry.

## FOR USE ON 5 - 55 GALLON CONTAINERS OR REFILLS

(For containers greater than 1 gallon)

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

#### PESTICIDE STORAGE

Store in area inaccessible to children or persons unfamiliar with product uses. Do not store on side.

Avoid creasing or impacting walls

#### PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray or mixture of rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration or if allowed by state and local authorities by burning. If burned, stay out of smoke. Metal Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning or dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

**If container is 5 gallons or more, the following statement must be used:**

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

**This product is effective against the following microorganisms:**  
Contact time is 10 minutes unless noted as below.

**Isolates From AIDS Patients**

- 1 *Aspergillus niger* (FOR CALIFORNIA USE ONLY: Contact time is 20 minutes)
- 2 *Candida albicans*
- 3 *Cryptococcus neoformans*
- 4 *Pseudomonas aeruginosa*
- 5 *Staphylococcus aureus*
- 6 *Streptococcus pneumoniae*

**Gram Positive Clinical Isolates**

- 7 *Enterococcus faecalis*
- 8 *Micrococcus luteus*
- 9 *Staphylococcus aureus*
- 10 *Staphylococcus aureus* (Toxic shock)
- 11 *Staphylococcus epidermidis*
- 12 *Staphylococcus saprophyticus*
- 13 *Streptococcus haemolyticus*

**Gram Negative Clinical Isolates**

- 14 *Acinetobacter calcoaceticus* var. *anitratus*
- 15 *Acinetobacter calcoaceticus* var. *lwoffii*
- 16 *Bordetella bronchiseptica*
- 17 *Brevundimonas dimunita*
- 18 *Burkholderia cepacia*
- 19 *Enterobacter agglomerans*
- 20 *Enterobacter cloacae*
- 21 *Enterobacter gergoviae*
- 22 *Enterobacter liquefaciens*
- 23 *Escherichia coli* (Urinary)
- 24 *Escherichia coli* (Wound)
- 25 *Flavobacterium meningosepticum*
- 26 *Hafnia alvei*
- 27 *Klebsiella oxytoca*
- 28 *Klebsiella pneumoniae*
- 29 *Morganella morganii*
- 30 *Proteus mirabilis*
- 31 *Proteus vulgaris*
- 32 *Pseudomonas aeruginosa*
- 33 *Pseudomonas fluorescens*
- 34 *Pseudomonas pseudomallei*
- 35 *Pseudomonas putida*
- 36 *Pseudomonas stutzeri*
- 37 *Serratia marcescens*
- 38 *Sphingomonas paucimobilis*

# PUBLIC PLACES ORGANISMS INSERT

## Other Bacteria

- 39 *Actinobacillus pleuropneumoniae*
- 40 *Actinomyces pyogenes*
- 41 *Bordetella bronchiseptica*
- 42 *Burkholderia picketti*
- 43 *Campylobacter jejuni*
- 44 *Corynebacterium pseudotuberculosis*
- 45 *Escherichia coli* strain 0157:H7
- 46 *Escherichia vulneris*
- 47 *Legionella pneumophila*
- 48 *Listeria monocytogenes*
- 49 *Neisseria gonorrhoeae*
- 50 *Pasteurella haemolytica*
- 51 *Pseudomonas aeruginosa*
- 52 *Rhodococcus equi*
- 53 *Salmonella choleraesuis*
- 54 *Salmonella typhi*
- 55 *Staphylococcus auricularis*
- 56 *Staphylococcus capitis*
- 57 *Staphylococcus hominis*
- 58 *Staphylococcus simulans*
- 59 *Streptococcus equi* var. *equi*
- 60 *Streptococcus equi* var. *zooepidermicus*
- 61 *Streptococcus pyogenes*
- 62 *Treponema pallidum* (ATCC # 27087)
- 63 *Yersinia enterocolitica*

## Pathogenic Fungi

- 64 *Trichophyton mentagrophytes*

## Environmental Fungi

- 65 *Aspergillus candidus*
- 66 *Aspergillus niger* (FOR CALIFORNIA USE ONLY: Contact time is 20 minutes)
- 67 *Penicillium chermesinum*
- 68 *Penicillium oxalicum*
- 69 *Penicillium spinulosum*
- 70 *Ulocladium* sp.

## Antibiotic Resistant Gram Positive Rods

- 71 *Enterococcus faecalis* (Vancomycin Resistant)
- 72 *Staphylococcus aureus* (Methicillin Resistant)

### **Human Viruses**

- 73 Cytomegalovirus
- 74 HBV (Hepatitis B Virus)
- 75 Herpes Simplex type 1 Virus
- 76 Herpes Simplex type 2 Virus
- 77 HIV-1 (AIDS Virus, human immunodeficiency virus Type 1)
- 78 Influenza A2-Asian Virus
- 79 Influenza A/Brazil Virus
- 80 Influenza B Virus (Allen strain)
- 81 Influenza C Virus (Taylor strain)
- 82 Measles Virus
- 83 Parainfluenza type 1
- 84 Poliovirus type 1 (Chat strain) **30 minutes contact time**

### **Non-Human Viruses**

- 85 Avian Influenza/Turkey/Wisconsin Virus
- 86 Equine Influenza
- 87 Feline Calicivirus
- 88 Feline Infectious Peritonitis
- 89 Infectious Bovine Rhinotracheitis (IBR)
- 90 Newcastle Disease Virus
- 91 Porcine Parvovirus
- 92 Pseudorabies Virus
- 93 Transmissible Gastroenteritis (TGE)
- 94 T1 bacteriophage
- 95 T4 bacteriophage